

# CBCS SCHEME

18MAT21

## Second Semester B.E. Degree Examination, Aug./Sept.2020 **Advanced Calculus and Numerical Methods**

Time: 3 hrs.

Max. Marks:100

Note: Answer any FIVE full questions, choosing ONE full question from each module.

a. Find the angle between the surfaces  $x^2 + y^2 - z^2 = 4$  and  $z = x^2 + y^2 - 13$  at (2, 1, 2). (06 Marks)

b. If  $\overrightarrow{F} = \nabla(xy^3z^2)$ , find div  $\overrightarrow{F}$  and curl  $\overrightarrow{F}$  at (1,-1,1). (07 Marks)

c. Find the value of the constant a such that the vector field

 $\vec{F} = (axy - z^3) \hat{i} + (a-2)x^2 \hat{j} + (1-a)xz^2 \hat{k}$ 

is irrotational and hence find a scalar function  $\phi$  such that  $\vec{F} = \nabla \phi$ . (07 Marks)

2 a. If  $\vec{F} = (3x^2 + 6y)i - 14yzj + 20xz^2k$ , evaluate  $\int \vec{F} \cdot d\vec{r}$  from (0, 0, 0) to (1, 1, 1) along the curve

given by x = t,  $y = t^2$  and  $z = t^3$ . (06 Marks) b. Use Green's theorem to find the area between the parabolas  $x^2 = 4y$  and  $y^2 = 4x$ . (07 Marks)

c. If  $\vec{F} = 2xyi + yz^2j + xzk$  and s is the rectangular parallelopiped bounded by x = 0, y = 0, z = 0and x = 2, y = 1, z = 3. Find the flux across S. (07 Marks)

Module-2

a. Solve  $(D^2 + 3D + 2)y = 4 \cos^2 x$ .

(06 Marks)

b. Solve  $(D^2 + 1)y = \sec x \tan x$ , by the method of variation of parameter. c. Solve  $x^2y'' + xy' + 9 = 3x^2 + \sin(3\log x)$ .

(07 Marks)

(07 Marks)

OR

a. Solve  $y'' + 2y' + y = 2x + x^2$ .

(06 Marks)

b. Solve  $(2x+1)^2y'' - 6(2x+1)y' + 16y = 8(2x+1)^2$ .

(07 Marks)

c. The current i and the charge q in a series circuit containing on inductance L, capacitance C, emf E satisfy the differential equation:  $L\frac{di}{dt} + \frac{q}{c} = E$ ;  $i = \frac{dq}{dt}$ . Express q and i interms of t, given that L, C, E are constants and the value of i, q are both zero initially. (07 Marks)

Module-3

a. Form the partial differential equation by eliminating the arbitrary function from  $\phi(xy + z^2, x + y + z) = 0.$ (06 Marks)

b. Solve  $\frac{\partial^2 z}{\partial x \partial y} = \sin x \sin y$  for which  $\frac{\partial z}{\partial y} = -2 \sin y$  when x = 0 and z = 0 if  $y = (2n+1)\frac{\pi}{2}$ . (07 Marks)

Derive one dimensional wave equation in the standard form  $\frac{\partial^2 \mathbf{u}}{\partial \mathbf{r}^2} = \mathbf{c}^2 \frac{\partial^2 \mathbf{u}}{\partial \mathbf{v}^2}$ . (07 Marks)

### OR

- a. Form the partial differential equation by eliminating the arbitrary function form  $f\left(\frac{xy}{z},z\right)=0$ . (06 Marks)
  - b. Solve  $\frac{\partial^2 z}{\partial y^2} = z$ , given that when y = 0,  $z = e^x$  and  $\frac{\partial z}{\partial v} = e^{-x}$ . (07 Marks)
  - c. Find all possible solutions of one dimensional heat equation  $\frac{\partial u}{\partial t} = C^2 \frac{\partial^2 u}{\partial x^2}$  using the method (07 Marks) of separation of variables.

- Test for convergence of the series  $\sum_{n=1}^{\infty} \frac{(n!)^2}{(2n)!} x^n, (x > 0).$ (06 Marks)
  - Solve the Bessel's differential equation leading to  $J_n(x)$ . (07 Marks)
  - Express  $f(x) = x^4 + 3x^3 x^2 + 5x 2$  in terms of Legendre's polynomials. (07 Marks)

- Test for convergence of the series  $\sum_{n=1}^{\infty} \frac{n^2}{2^n}$ . (06 Marks)
  - If  $\alpha$  and  $\beta$  are two distinct roots fo  $J_n(x)=0$ . Prove that  $\int xJ_n(\alpha x)J_n(\beta x)dx=0$ . If  $\alpha\neq\beta$ .
  - (07 Marks) Express  $f(x) = x^3 + 2x^2 - x - 3$  in terms of Legendre's polynomials. (07 Marks)

- Find the real root of the equation :  $\frac{\text{Module-5}}{x^3 2x 5} = 0$  using Regula Falsi method, correct to (06 Marks) three decimal places.
  - b. Use Lagrange's formula, find the interpolating polynomial that approximates the function described by the following data:

X	0	1	2	5
f(x)	2	3	12	147

c. Evaluate  $\int_{0}^{1} \frac{x dx}{1 + x^2}$  by Weddle's rule, taking seven ordinates and hence find log  $e^2$ .

- Find the real root of the equation  $xe^x 2 = 0$  using Newton Raphson method correct to 10 three decimal places.
  - b. Use Newton's divided difference formula to find f(4) given the data:

1	x	0	2	3	6
	f(x)	-4	2	14	158

c. Use Simpson's  $\frac{3}{8}$  rule to evaluate  $\int_{0}^{4} e^{1/x} dx$ .