

# CBCS SCHEME

USN

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

18MAT31

## Third Semester B.E. Degree Examination, July/August 2021 Transform Calculus, Fourier Series and Numerical Techniques

Time: 3 hrs.

Max. Marks:100

**Note: Answer any FIVE full questions.**

- 1 a. Find  $L[t e^{-2t} \sin 4t]$ . (06 Marks)
  - b. A periodic function of period  $2\pi/\omega$  is defined by  $f(t) = \begin{cases} E \sin \omega t, & 0 \leq t < \pi/\omega \\ 0, & \pi/\omega \leq t < 2\pi/\omega \end{cases}$ . Where E and  $\omega$  are constants. (07 Marks)
  - c. Solve :  $y''(t) + k^2 y(t) = 0$ ;  $y(0) = 0$  and  $y'(0) = 1$  by Laplace transformation. (07 Marks)
  - 2 a. Find : i)  $L^{-1}\left\{\frac{s^2 - 3s + 4}{s^3}\right\}$  ii)  $L^{-1}\left[\text{Cot}^{-1}\left(\frac{S}{2}\right)\right]$ . (06 Marks)
  - b. Find the inverse Laplace transform of  $\frac{1}{(s-1)(s^2+1)}$  by using convolution theorem. (07 Marks)
  - c. Express the following function in terms of Heaviside step function and hence find its Laplace transform where  $f(t) = \begin{cases} t^2, & 0 < t \leq 2 \\ 4t, & t > 2 \end{cases}$ . (07 Marks)
  - 3 a. Expand  $f(x) = x(2\pi - x)$  as a Fourier series in  $[0, 2\pi]$ . (06 Marks)
  - b. Obtain Fourier series for the function  $f(x)$  given by  $f(x) = \begin{cases} 1 + \frac{2x}{\pi}, & -\pi \leq x \leq 0 \\ 1 - \frac{2x}{\pi}, & 0 \leq x \leq \pi \end{cases}$ . (07 Marks)
  - c. Find the half range sine series of  $f(x) = \frac{e^{ax}}{\sinh a \pi}$  in  $(0, \pi)$ . (07 Marks)
  - 4 a. Find the Fourier series expansion of  $f(x)$  given by  $f(x) = \begin{cases} 1 & 0 < x < 1 \\ 2 & 1 < x < 3 \end{cases}$ . (06 Marks)
  - b. Find the half range sine series for  $x^2$  in  $(0, \pi)$ . (07 Marks)
  - c. The values of  $x$  and the corresponding values of  $f(x)$  over a period T are given below. Show that  $f(x) = 0.75 + 0.37 \cos \theta + 1.004 \sin \theta$  where  $\theta = \frac{2\pi x}{T}$ . (07 Marks)
- |      |      |      |      |      |       |       |      |
|------|------|------|------|------|-------|-------|------|
| x    | 0    | T/6  | T/3  | T/2  | 2T/3  | 5T/6  | T    |
| f(x) | 1.98 | 1.30 | 1.05 | 1.30 | -0.88 | -0.25 | 1.98 |
- 5 a. State: i) Initial and final value theorems ii) Find the Z -transform of  $\cos\left(\frac{n\pi}{2} + \frac{\pi}{4}\right)$ . (06 Marks)
  - b. Find the complex Fourier transform of the function  $f(x) = \begin{cases} 1 & \text{for } |x| \leq a \\ 0 & \text{for } |x| > a \end{cases}$   
Hence evaluate  $\int_0^{\infty} \left(\frac{\sin x}{x}\right) dx$ . (07 Marks)
  - c. Compute the inverse Z-transform of  $\frac{3z^2 + 2z}{(5z-1)(5z+2)}$ . (07 Marks)

- 6 a. Find the Fourier cosine transform of  $f(x) = \begin{cases} x, & 0 < x < 2 \\ 0, & \text{else where} \end{cases}$ . (06 Marks)
- b. Find the Z-transform of  $2n + \sin \frac{n\pi}{4} + 1$ . (07 Marks)
- c. Solve the difference equation :  $u_{n+2} - 3u_{n+1} + 2u_n = 0$ , with  $u_0 = 0$  and  $u_1 = -1$ . (07 Marks)
- 7 a. Find by Taylor's series method the value of  $y$  at  $x = 0.1$  to five places of decimals from  $\frac{dy}{dx} = x^2y - 1, y(0) = 1$ . (06 Marks)
- b. Use fourth order Runge-Kutta method to solve  $(x + y)\frac{dy}{dx} = 1, y(0.4) = 1$  at  $x = 0.5$  correct to four decimal places. (07 Marks)
- c. If  $\frac{dy}{dx} = 2e^x - y, y(0) = 2, y(0.1) = 2.010, y(0.2) = 2.040$  and  $y(0.3) = 2.090$ , find  $y(0.4)$  correct to four decimal places by using Milne's predictor - corrector method and applying corrector formula twice. (07 Marks)
- 8 a. Using modified Euler's formula compute  $y(1.1)$  correct to five decimal places given that  $\frac{dy}{dx} + \frac{y}{x} = \frac{1}{x^2}$  and  $y = 1$  at  $x = 1$ . [taking  $h = 0.1$ ]. (06 Marks)
- b. Employ Taylor's series method to find  $y$  at  $x = 0.1$  and  $0.2$  correct to four places of decimal. Given  $\frac{dy}{dx} - 2y = 3e^x, y(0) = 0$ . (07 Marks)
- c. Solve the differential equation  $y' + y + xy^2 = 0$  with the initial values of  $y : y_0 = 1, y_1 = 0.9008, y_2 = 0.8066, y_3 = 0.722$  corresponding to the values of  $x : x_0 = 0, x_1 = 0.1, x_2 = 0.2, x_3 = 0.3$  by computing the value of  $y$  corresponding to  $x = 0.4$  applying Adams - Bashforth predictor and corrector formula. (07 Marks)
- 9 a. Given  $y'' - xy' - y = 0$  with the initial conditions  $y(0) = 1, y'(0) = 0$ , compute  $y(0.2)$  using fourth order Runge-Kutta method. (06 Marks)
- b. Derive Euler's equation in the standard form  $\frac{\partial f}{\partial y} - \frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) = 0$ . (07 Marks)
- c. A heavy cable hangs freely under gravity between two fixed points. Show that the shape of the cable is a catenary. (07 Marks)
- 10 a. Apply Milne's method to compute  $y(0.8)$  given that  $y'' = 1 - 2yy'$  and the following table of initial values. (07 Marks)
- |    |   |        |        |        |
|----|---|--------|--------|--------|
| x  | 0 | 0.2    | 0.4    | 0.6    |
| y  | 0 | 0.02   | 0.0795 | 0.1762 |
| y' | 0 | 0.1996 | 0.3937 | 0.5689 |
- b. Prove that the geodesics on a plane are straight line. (06 Marks)
- c. Find the extremal of the functional :  $\int_{x_0}^{x_1} (y^2 + y'^2 - 2y \sin x) dx$ . (07 Marks)

\*\*\*\*\*