

Synthesis, Characterization, Anti-Microbial and DNA Binding Studies on Mn(II) and Zn(II) Metal Complexes Containing Mixed Ligands

B.Sreekanth^{1*}, T. K. Vishnuvardhan², Veena V Pillai¹, P. Murali Krishna³,
N. B.Gopal Reddy³, Jagan Mohan Reddy¹, V.Sridhara⁴

¹Department of Biotechnology, ²Department of Chemistry, Acharya Institute of Technology, Bangalore -560 107,India.

³Department of Chemistry, M.S Ramaiah Institute of Technology, Bangalore-560 054,India.

⁴Department of Biotechnology, New Horizon College of Engineering, Bangalore -560 103,India.

*Corres. Author:sreekanthb@acharya.ac.in
Mobile No: +91-7829174813

Abstract: The DNA binding ability of the novel complexes $Mn(L_1)_2L_2(PF_6)_2$ [Complex (1) and $Zn(L_1)_2L_2(PF_6)_2$ [Complex (2)] containing bioactive mixed ligand of type $L_1=8$ -Hydroxyquinoline and $L_2=1,10$ -Phenanthroline have been isolated and characterized by analytical and spectral methods. The intrinsic binding constant K_b has been estimated at room temperature. The binding constant of the complexes in 5 mM Tris-HCl/50 mM NaCl buffer at pH 7.2, are $13.10 \times 10^6 M^{-1}$ and $17.41 \times 10^6 M^{-1}$ for complex (1) and (2) respectively. The absorption spectra indicate that the complexes intercalate between the base pairs of the CT-DNA tightly. The synthesized compounds have been tested against microorganisms such as Gram positive bacteria (*Staphylococcus aureus* and *Proteus vulgaris*), Gram-negative bacteria (*Escherichia coli* and *Pseudomonas aeruginosa*) and fungus (*Aspergillus niger*). A comparative study of the Minimum Inhibitory Concentration (MIC) values of the ligands and their complexes indicates that the complexes exhibit moderate antimicrobial activity than the free ligand and control.

Keywords: Manganese and Zinc complexes, DNA Binding and Antimicrobial studies.

INTRODUCTION

Organometallic chemistry has been developed, in the last four decades, to be the largest and important branch as a link connecting the fields of organic and inorganic chemistry. One of the major applications of the transition metal complexes is its medical testing as antibacterial and antitumor agents aiming toward the discovery of an effective and safe therapeutic regimen for the treatment of bacterial infections and cancers. In addition, a great many Schiff base complexes with metals have also provoked wide interest because they possess a diverse spectrum of biological and pharmaceutical activities, including antitumor and antioxidative activities. Moreover, mixed-ligand complexes are observed in biological systems or in the intermediate

chemical reactions with metal ions, which are important to understand the respective chemistry. Investigations concerning 1,10-phenanthroline mixed-ligand chelate systems would provide toward understanding the driving forces that led to the formation of such mixed ligand complexes¹. Moreover, Cu(II) and Zn(II) complexes with oxygen, sulphur and nitrogen containing ligands are the subject of intensive biological evaluation in the search for less toxic and more selective anticancer therapies^{2,3}.

The study of mixed ligand-complex formation is relevant in the field of analytical chemistry, where the use of mixed ligand complexes allows the development of methods with increased selectivity, sensitivity and has also great importance in the field of biological and environmental chemistry^{4,5}.

The study of mixed ligand-complex formation is relevant in the field of analytical chemistry, where the use of mixed ligand complexes allows the development of methods with increased selectivity, sensitivity and has also great importance in the field of biological and environmental chemistry^{6,7}. The effect of the presence of methyl substituent in the phenyl rings of aromatic Schiff bases on their antimicrobial activity has been reported⁸. Although many Schiff bases are known to be active against a wide range of micro-organisms, for example; *E.coli*, *B.subtilis*, and *S.aureus* and fungi *R.stolonifer* and *C.ablican*, antibacterial activity has been studied more than antifungal activity, since bacteria are usually more resistant to antibiotics through biochemical and morphological modifications. Moreover, the incorporation of transition metal into Schiff bases enhances the biological activity of the ligand and decreases the cytotoxic effects of both the metal ion and ligand on the host⁹.

However the literature survey that the mixed ligands of 8-Hydroxyquinoline and 1-10 Phenanthroline derivatives with their transition metal complexes have not been reported and studied so far. Hence the present study aims for the Synthesis, Characterization, Antimicrobial and DNA binding studies of Manganese and Zinc complexes containing mixed ligands 8-hydroxyquinoline and 1-10 Phenanthroline.

EXPERIMENTAL

All reagents and solvents were of AR grade, solvents were purified and used. Metal salts such as Manganese Sulphate and Zinc Sulphate were purchased from Karnataka fine chemicals. Ligands such as 8-Hydroxyquinoline and 1,10-Phenanthroline was purchased from Himedia chemicals Ltd. (Bangalore).


SYNTHESIS OF METAL COMPLEXES

Synthesis of Manganese complex ($Mn(L_1)_2L_2$) (PF_6)₂[Complex (1)]

Manganese sulphate (0.169 g, 1mmol) 8-hydroxyquinoline L_1 (0.29 g, 2mmol) 1,10-phenanthroline L_2 (0.234g, 1mmol) were dissolved in ethanolic solution and refluxed on the water bath for 4 hours. The contents were cooled to obtain precipitate. The complex was filtered and dried under vacuum before being recrystallized in acetone.

Synthesis of Zinc complex ($Zn(L_1)_2L_2$) (PF_6)₂[Complex (2)]

Zinc sulphate (0.287 g, 1mmol) 8-hydroxyquinoline L_1 (0.29 g, 2mmol) 1,10-phenanthroline L_2 (0.234g, 1mmol) were dissolved in ethanolic solution and refluxed on the water bath for 4 hours. The contents were cooled to obtain precipitate. The complex was filtered and dried under vacuum before being recrystallized in acetone.



Where M= Mn or Zn

Figure 1: Proposed structure of the metal complexes

SPECTRAL MEASUREMENTS

IR spectra were recorded with Shimadzu model FT-IR spectrophotometer by using KBr pellets. UV-visible absorption spectra were recorded using ELICO model SL-159 UV-Vis Spectrophotometer at room temperature.

DNA BINDING STUDIES

The concentration of CT-DNA per nucleotide [C(p)] was measured by using its known extinction coefficient at 260 nm ($6600 \text{ M}^{-1} \text{ cm}^{-1}$)¹⁰. TrisHCl-buffer [5mM Tris(hydroxymethyl) amino methane, pH 7.2, 50 mM NaCl] was used for the absorption, viscosity, and thermal denaturation experiments.

Absorption titration experiments were carried out by varying the DNA concentration (0-100 μM) and maintaining the metal complex concentration constant. Absorption spectra were recorded after each successive addition of DNA and equilibration (approximately 10 minutes). For both (1) and (2), the observed data were then fitted into Equation (1) to obtain the intrinsic binding constant K_b ¹¹.

$$[\text{DNA}] / (a - f) = [\text{DNA}] / (b - f) + 1/K_b (a - f),$$

where a , b , and f are the apparent, bound, and free metal complex extinction coefficients, respectively, at 263 nm for (1) and (2). A plot of $[\text{DNA}] / (b - f)$ versus $[\text{DNA}]$ gave a slope of $1/(b - f)$ and a y intercept equal to $1/K_b (b - f)$, where K_b is ratio of the slope to the y intercept.

ANTIMICROBIAL STUDIES

The ligands and its complexes were investigated for anti-bacterial and anti-fungal properties. Gram positive bacteria (*Staphylococcus aureus* and *Proteus vulgaris*), Gram-negative bacteria (*Escherichia coli* and *Pseudomonas aeruginosa*) and one fungus (*Aspergillus niger*) were used in this study to assess their antimicrobial properties. The tested compounds were dissolved in DMSO and the solutions were serially diluted in order to find the MIC values. The antibiotic Chloramphenicol was used as standard reference in the case of Gram-negative bacteria, Amikacin was used as standard reference in case of Gram-positive bacteria and Clotrimazole was used as standard anti-fungal reference. The solvent DMSO was used as negative control. In a typical procedure, a well was made in the Muller Hinton agar medium inoculated with microorganisms. The well was filled with the test solution using a micropipette and the plates were incubated at 37 °C at 72 h for fungi and 24 h for bacteria. During this period, the test solution diffused and affected the growth of the inoculated fungi and bacteria. Activity was determined by measuring the diameter of the zone of inhibition (mm).

RESULTS

CHARACTERIZATION OF COMPLEXES

The elemental analysis data are agreed with the theoretical values within the limit of experimental error. These complexes are soluble in DMF, DMSO and in buffer (pH 7.2) solution.

The IR spectra of complexes were recorded in the range of 4000-400 cm^{-1} on KBr pellets. The spectra of the ligands 8-Hydroxyquinoline and 1,10-Phenanthroline showed bands at 1577 cm^{-1} and 1597 cm^{-1} assigned to $\nu\text{C}=\text{N}$ aromatic hydrocarbon, 3045 cm^{-1} and 3024 cm^{-1} assigned to $\nu\text{C}-\text{H}$ group and 3600 cm^{-1} and 3800 cm^{-1} assigned to $\nu\text{O}-\text{H}$ group. The spectra of both the complexes showed a peak in the range 1510 cm^{-1} to 1630 cm^{-1} for $\nu\text{C}=\text{N}$ group are shifted slightly indicating that the coordination taken place through nitrogen atom.

ABSORPTION SPECTRAL STUDIES

The absorption spectra of complexes (1) and (2) in the absence and presence of CT-DNA are given in Figures 2 and 3, respectively. Figure 2 and 3 depicts a well-resolved band at 263 nm for complex (1) and complex (2) with increasing DNA concentrations (0-100 μM). The result shows that the absorbance (hypochromism) decreased by the successive addition of CT-DNA to the complex solution. The hypochromism observed for the bands of complexes (1) and (2) is accompanied by a small bathochromic shift of 2 and 1 nm, as shown in Figures 2 and 3, respectively. The hypochromism and bathochromic shift observed for the complexes suggest that binding is

in intercalative mode. To compare quantitatively the DNA binding strengths of these complexes, the intrinsic DNA binding constants K_b are determined from the decay of the absorbance at 263 nm for complex (1) and complex (2) with increasing concentrations of DNA by using equation (1) given in our previous article^{11,18}. The observed K_b values for complexes (1) and (2) are equal to the classical intercalators bound to CT-DNA. The K_b values for complexes (1) and (2) are $13.10 \times 10^6 \text{ M}^{-1}$ and $17.41 \times 10^6 \text{ M}^{-1}$, respectively. Thus, it is obvious that the present complexes are involved in intercalative interactions with CT-DNA.




Figure 2: Absorption spectra of complex (1) in Tris-HCl buffer upon addition of DNA. $[\text{Mn}] = 0.5 \mu\text{M}$, $[\text{DNA}] = 0\text{--}100 \mu\text{M}$. Arrow shows the absorbance changing upon the increase of DNA concentration.

Inset: The plot of $[\text{DNA}]/(a - f)$ versus $[\text{DNA}]$ for the titration of DNA with Mn(II) complex.




Figure 3: Absorption spectra of complex (2) in Tris-HCl buffer upon addition of DNA. $[\text{Zn}] = 0.5 \mu\text{M}$, $[\text{DNA}] = 0\text{--}100 \mu\text{M}$. Arrow shows the absorbance changing upon the increase of DNA concentration.

Inset: The plot of $[\text{DNA}]/(a - f)$ versus $[\text{DNA}]$ for the titration of DNA with Zn(II) complex.

ANTIMICROBIAL STUDIES

The in vitro antimicrobial activity of the compounds was tested against the bacterial species including *Staphylococcus aureus*, *Proteus vulgaris*, *Escherichia coli* and *Pseudomonas aeruginosa* and the fungus *Aspergillus niger* by well diffusion method. These complexes are inhibiting Gram-positive and Gram negative bacterial strains. The importance of this unique property of the investigated Schiff base complexes lies in the fact that, it can be applied safely in the treatment of infections and some common diseases e.g. Septicaemia, Gastroenteritis, Urinary tract infections and hospital acquired infections¹⁹. The ligand and their complexes have been tested for in vitro growth inhibitory activity against gram-positive microbe *Staphylococcus aureus*, *Proteus*

vulgaris and gram-negative microbe's *Escherichia coli*, *Pseudomonas aeruginosa* by using well-diffusion method. As the test solution concentration increases, the biological activity also increases. The Minimum Inhibitory Concentration (MIC) values of the investigated compounds are summarized in Table 1. From the table, the observed MIC values indicate that the complexes have higher antibacterial activity. But the complexes shows no inhibition against fungi. (*Aspergillus niger*). The metal complexes Mn(II) and Zn(II) have higher antibacterial activity than the ligand are shown in Figs. 4(a), 4(b) and 4(c). The increase in antimicrobial activity is due to the faster diffusion of metal complexes as a whole.




Fig4 (a): Activity of *E. coli* against Ligands and Complex (1) and (2)




Fig4 (b): Activity of *P. aeruginosa* against Ligands and Complex (1) and (2)




Fig4(c): Activity of *S. aureus* against Ligands and Complex (1) and (2)




Fig4 (d): Activity of *Proteus vulgaris* against Ligands and Complex (1) and (2)

Table 1: Antimicrobial activity of the ligand and metal complexes

Sl.No	Extract	Conc (mg/ml)	<i>E.coli</i> (mm)	<i>P. aeruginosa</i> (mm)	<i>Staphylococcus aureus</i> (mm)	<i>Proteus vulgaris</i> (mm)	<i>A.niger</i> (mm)
1	L ₁	0.5	0	8	0	0	0
		1	0	15	0	0	0
		1.5	0	16	0	0	0
		2	0	17	12	0	0
		2.5	0	19	15	0	0
2	L ₂	0.5	0	0	0	12	0
		1	0	10	10	15	0
		1.5	15	12	16	16	0
		2	22	15	19	21	0
		2.5	24	22	23	22	0
3	Mn(L ₁) ² L ₂	0.5	0	0	10	0	0
		1	9	0	16	10	0
		1.5	13	9	17	14	0
		2	15	10	18	16	0
		2.5	20	16	22	20	0
4	Zn(L ₁) ² L ₂	0.5	0	0	0	0	0
		1	0	0	14	15	0
		1.5	9	10	18	20	0
		2	17	16	21	21	0
		2.5	21	20	20	22	0
5	Antibiotic	0.5	21	24	21	21	0
		1	21	22	20	22	0
		1.5	22	22	21	22	0
		2	24	24	24	24	0
		2.5	22	24	25	24	0
6	DMSO	0.5	0	0	0	0	0
		1	0	0	0	0	0
		1.5	0	0	0	0	0
		2	0	0	0	0	0
		2.5	0	0	0	0	0

Less than 10mm-- Inactive; Less than 10-15mm--Weakly active; Less than 15-20mm--Moderately active;More than 20mm—Highly active

Standard: Antibiotic Chloramphenicol for Gram negative bacteria, Amikacin for Gram-positive bacteria, Clotrimazole for fungi

Control: DMSO

CONCLUSION

In conclusion, we have synthesized and characterized two new complexes of the type $[\text{Mn}(\text{L}_1)_2\text{L}_2](\text{PF}_6)_2$ (**1**) and $[\text{Zn}(\text{L}_1)_2\text{L}_2](\text{PF}_6)_2$ (**2**). Interactions of the new complexes with (double stranded) DNA were investigated by absorption spectra, viscosity and thermal denaturation studies. From the experimental results, it was confirmed that the complexes bound with the double stranded DNA with binding constant $K_b = 13.10 \times 10^6 \text{ M}^{-1}$ for (**1**) and $K_b = 17.41 \times 10^4 \text{ M}^{-1}$ for (**2**), respectively. The metal complexes have higher antimicrobial activity than the free ligand.

ACKNOWLEDGEMENTS

We would like to acknowledge the management and staff of Acharya Institute of Technology, Bangalore and M.S Ramaiah Institute of Technology, Bangalore.

REFERENCES

1. Raman, N. and Sobha, S. Synthesis, Characterization, DNA interaction and antimicrobial screening of isatin-based polypyridyl mixed-ligand Cu(II) and Zn(II) complexes, *J. Serb.Chem. Soc.*, 2010, 75,1-16.
2. Jakupec, M.J, Galanski, M, and Keppler, B.K., Tumour-inhibiting platinum complexes state of the art and future perspectives, *Rev. Physiol. Biochem. Pharmacol*, 2003,146,1–53.
3. Gaballa, A.S, Asker, M.S, Barakat, A.S, and Teleb, S.M., Synthesis, characterization and biological activity of some platinum(II) complexes with Schiff bases derived from salicylaldehyde, 2-furaldehyde and phenylenediamine, *Specrochim. Acta.*,2007, 67A, 114-121.
4. Keskioglu, E, Gunduzalp, A.B, Cete, S, Hamurcu, F, and Erk, B., Cr(III), Fe(III) and Co(III) complexes of tetradentate (ONNO) Schiff base ligands: Synthesis, characterization, properties and biological activity, *Spectrochim. Acta.*,2008, 70, 634-640.
5. Reichmann, M.E, Rice, S.A, Thomas, C.A and Doty, P., A further examination of the molecular weight and size of deoxypentose nucleic acid,*J. Am. Chem. Soc.*, 1954, 76, 3047–3053.
6. Pyle, A. M, Barton,J. K, and Lippard, S.J.,Oxidative DNA damage through long-range electron transfer,*J. Am. Chem. Soc.*, 1992, 114, 2303-2311.
7. Tullius, T. D., *Metal-DNA Chemistry*. ACS Symposium Series, American Chemical Society, Washington, DC, 2008,402.
8. Shivarama Holla, B., Mahalinga, M., Sithambaram Karthikeyan, M., Poojary, B., Mohammed Akberlib, P. and Suchetha Kumar, N., Synthesis, characterization and antimicrobial activity of some substituted 1,2,3-triazoles, *Eur. J. Med. Chem.*, 2005, 40, 1173-1178.
9. Ameerunisha Begum, M.S, Saha, S, Nethaji, M. and Chakravarty, A.R., Iron(III) Schiff base complexes of arginine and lysine as netropsin mimics showing AT-selective DNA binding and photonuclease activity, *J. Inorg. Biochem.*, 2010, 104, 477-484.
10. Reichmann, M.E., Rice, S.A., Thomas, C.A.,and Doty, P., A further examination of the molecular weight and size of deoxypentose nucleic acid,*J. Am. Chem. Soc.* 1954, 76, 3047–3053.
11. Wolfe, A., Shimer, G.H.,and Meehan, T., Polycyclic aromatic hydrocarbons physically intercalate into duplex regions of denatured DNA, *Biochemistry*,1987, 26, 6392–6396.
12. Sreekanth, B., Krishnamurthy, G., Bhojya Naik, H.S., Vishnuvardhan, T.K., Vinaykumar, B., and Sharath, N., Synthesis, DNAbinding, and oxidative cleavage studies of Fe(II) andCo(III) complexes containing bioactive ligands,*Nucleosides Nucleotides Nucleic Acids*,2011, 30, 83–96.
13. Sreekanth, B., Krishnamurthy, G., Bhojya Naik, H. S., Vishnuvardhan, T.K.,and Prabhakar, M.C., Fe(II) complexes containing bioactive ligands: Synthesis, DNA binding, and cleavage studies. *Synth. React.Inorg. Met-Org. Nano-Met. Chem.*, 2010, 40, 955–962.
14. Sreekanth, B., Krishnamurthy, G., Bhojya Naik, H. S., Vishnuvardhan, T. K., Lokesh, M.R., and Shashikumar, N. D.,DNA Interaction Studies of Cu (II) and Mn (II) Complexes containing Mixed Ligand of 1, 10-phenanthroline and 2-hydroxy-4-methyl-1, 8-naphthyridine,*Res. J. of Pharm., Biol. and Chem. Sci.*,2011, 2, 201-213
15. Sreekanth, B., Krishnamurthy, G., Bhojya Naik, H. S., Vishnuvardhan,T. K., Lokesh, M.R., and Shashikumar, N. D.,DNA Binding and Cleavage Studies of Fe(II) and Zn(II) Complexes containing

- Mixed Ligand of 1,10-phenanthroline and 2-hydroxy-4- methyl-1,8-naphthyridine. J. of Chem. and Pharm. Res., 2011, 3, 407-419.
16. Sreekanth, B., Krishnamurthy, G., Bhojya Naik, H.S.,and Vishnuvardhan,T. K.,Cu(II) and Mn(II) Complexes Containing Macroacyclic Ligand: Synthesis, DNA Binding, and Cleavage Studies, Nucleosides, Nucleotides and Nucliecacids, 2012, 31, 1-13.
 17. Sreekanth, B., Krishnamurthy, G., Bhojya Naik, H.S., Gopinath, S.M., Vishnuvardhan,T. K.,and Ismail Shareef.,Synthesis, Characterization, DNA Binding and Cleavage studies of Fe(III)and Zn(II) Complexes Containing Mixed Ligands. International Journal of Pharmacy & Technology, 2013, 4(3), 257-265.
 18. Sreekanth.B., and Veena V. Pillai.,DNA Binding and Anti-Microbial Studies of Ag(II) And Cu(II) Metal Complexes Containing Mixed Ligands of 1,10-Phenanthroline And 8-Hydroxyquinoline.International Journal of Pharma and Bio Sciences, 2013, (In Press).
 19. Natesan Mahalakshmi., and Dr.Rajavel,R., Binuclear Copper(II), Nickel(II) and Oxovanadium(IV) schiff base complexes bearing N₂O₂ donors and their DNA cleavage and antibacterial activity. International Journal of Pharmacy & Technology. 2010, 2, 1133-1157.
