

USN

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**Third Semester B.E. Degree Examination, Dec.2023/Jan.2024**  
**Transform Calculus Fourier Series & Numerical Techniques**

Time: 3 hrs.

Max. Marks: 100

*Note: Answer any FIVE full questions, choosing ONE full question from each module.*

**Module-1**

- 1
- a. Find the Laplace transform of, (i)  $e^{-3t} \sin 5t \cdot \cos 3t$  (ii)  $\frac{e^{at} - e^{bt}}{t}$ . (06 Marks)
- b. If a periodic function of period 'a' is defined by  $f(t) = \begin{cases} E, & \text{for } 0 < t < \frac{a}{2} \\ -E, & \text{for } \frac{a}{2} < t < a \end{cases}$  then show that  $L\{f(t)\} = \frac{E}{s} \tanh\left(\frac{as}{4}\right)$ . (07 Marks)
- c. Using convolution theorem find the inverse Laplace transform of  $\frac{s}{(s+2)(s^2+9)}$ . (07 Marks)

**OR**

- 2
- a. Express the function  $f(t) = \begin{cases} \cos t & \text{for } 0 < t < \pi \\ \cos 2t & \text{for } \pi < t < 2\pi \\ \cos 3t & \text{for } t > 2\pi \end{cases}$  in terms of unit step function and hence find its Laplace transform. (07 Marks)
- b. Find the inverse Laplace transform of  $\frac{2s^2 - 6s + 5}{s^3 - 6s^2 + 11s - 6}$ . (06 Marks)
- c. Solve the differential equation  $\frac{d^2y}{dt^2} + 4\frac{dy}{dt} + 4y = e^{-t}$  with  $y(0) = y'(0) = 0$  by using Laplace transform. (07 Marks)

**Module-2**

- 3
- a. Find a Fourier series to represent  $f(x) = |x|$  in  $-\pi \leq x \leq \pi$ . (06 Marks)
- b. Obtain the half-range cosine series for  $f(x) = x \sin x$  in  $(0, \pi)$  and hence show that  $\frac{\pi-2}{4} = \frac{1}{1.3} - \frac{1}{3.5} + \frac{1}{5.7} - \dots \infty$ . (07 Marks)
- c. Express y as a Fourier series up to second harmonics for the following data :

x:	0	$\frac{\pi}{3}$	$\frac{2\pi}{3}$	$\pi$	$\frac{4\pi}{3}$	$\frac{5\pi}{3}$	$2\pi$
y:	1	1.4	1.9	1.7	1.5	1.2	1.0

(07 Marks)

Important Note : 1. On completing your answers, compulsorily draw diagonal cross lines on the remaining blank pages.  
 2. Any revealing of identification, appeal to evaluator and /or equations written eg, 42+8 = 50, will be treated as malpractice.

OR

- 4 a. Obtain the Fourier series expansion for the function,  $f(x) = 2x - x^2$  in  $(0, 2)$ . (06 Marks)

- b. Find the half range sine series for the function,  $f(x) = \begin{cases} \frac{1}{4} - x & \text{for } 0 < x < \frac{1}{2} \\ x - \frac{3}{4} & \text{for } \frac{1}{2} < x < 1 \end{cases}$  (07 Marks)

- c. The following table gives the variation of periodic current over period :

t sec :	0	$\frac{T}{6}$	$\frac{T}{3}$	$\frac{T}{2}$	$\frac{2T}{3}$	$\frac{5T}{6}$	T
A (amp) :	1.98	1.30	1.05	1.30	-0.88	-0.25	1.98

Show that there is a direct current part of 0.75 amp in the variable current and obtain the amplitude of the first harmonic. (07 Marks)

**Module-3**

- 5 a. Find the Fourier transform of the function  $f(x) = \begin{cases} 1 - x^2 & \text{for } |x| \leq 1 \\ 0 & \text{for } |x| > 1 \end{cases}$ . Hence evaluate

$$\int_0^{\infty} \left( \frac{x \cos x - \sin x}{x^3} \right) dx. \quad (06 \text{ Marks})$$

- b. Find the Fourier sine and cosine transform of  $f(x) = \begin{cases} x & \text{if } 0 < x < 1 \\ 2 - x & \text{if } 1 < x < 2 \\ 0 & \text{otherwise} \end{cases}$ . (07 Marks)

- c. Find the z-transform of  $\cosh\left(n \frac{\pi}{2} + \theta\right)$ . (07 Marks)

OR

- 6 a. Find the Fourier sine transform of  $f(x) = e^{-ax}$ ,  $a > 0$ . (06 Marks)

- b. Find the inverse z transform of  $\frac{18z^2}{(2z-1)(4z+1)}$ . (07 Marks)

- c. Solve the difference equation  $u_{n+2} + 6u_{n+1} + 9u_n = z^n$  with  $u_0 = u_1 = 0$  using z-transform. (07 Marks)

**Module-4**

- 7 a. Classify the following partial differential equations :

(i)  $\frac{\partial^2 u}{\partial x^2} + 4 \frac{\partial^2 u}{\partial x \partial y} + 4 \frac{\partial^2 u}{\partial y^2} - \frac{\partial u}{\partial x} + 2 \frac{\partial u}{\partial y} = 0.$

(ii)  $x^2 \frac{\partial^2 u}{\partial x^2} + (1 - y^2) \frac{\partial^2 u}{\partial y^2} = 0, -\infty < x < \infty, -1 < y < 1.$

(iii)  $(1 + x^2) \frac{\partial^2 u}{\partial x^2} + (5 + 2x^2) \frac{\partial^2 u}{\partial x \partial t} + (4 + x^2) \frac{\partial^2 u}{\partial t^2} = 0.$

(iv)  $(x+1) \frac{\partial^2 u}{\partial x^2} - 2(x+2) \frac{\partial^2 u}{\partial x \partial y} + (x+3) \frac{\partial^2 u}{\partial y^2} = 0. \quad (10 \text{ Marks})$

- b. Evaluate the values at the mesh points for the equation  $u_{tt} = 16u_{xx}$  taking  $h = 1$  upto  $t = 1.25$ . The boundary conditions are  $u(0, t) = u(5, t) = 0$  and the initial conditions are  $u(x, 0) = x^2(5 - x)$  and  $u_t(x, 0) = 0$ . (10 Marks)

OR

- 8 a. Using Schmidt two-level formula to solve the equation  $\frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$  under the conditions,
- (i)  $u(0, t) = u(1, t) = 0 \quad t \geq 0$
- (ii)  $u(x, 0) = \sin \pi x, \quad 0 < x < 1$  by taking  $h = \frac{1}{4}$  and  $\alpha = \frac{1}{6}$  co. (10 Marks)
- b. Solve the two-dimensional Laplace equation  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$  at the interior mesh points of the square region and the values of  $u$  at the mesh points on the boundary are shown in Fig.Q8 (b).




Fig. Q8 (b)

(10 Marks)

**Module-5**

- 9 a. Using Runge-Kutta method of 4<sup>th</sup> order to solve the differential equation  $\frac{d^2 y}{dx^2} + 2x \frac{dy}{dx} - 4y = 0$  with  $y(0) = 0.2$  and  $y'(0) = 0.5$  for  $x = 0.1$ . Correct to four decimal places. (07 Marks)
- b. State and prove Euler's equation. (07 Marks)
- c. Find the extremal of the functional  $I = \int_0^{\frac{\pi}{2}} (y^2 - y'^2 - 2y \sin x) dx$  under the end conditions  $y(0) = 0, y\left(\frac{\pi}{2}\right) = 0$  (06 Marks)

OR

- 10 a. Apply Milne's method to compute  $y(0.3)$ . Given that  $\frac{d^2 y}{dx^2} = 1 - 2y \frac{dy}{dx}$  and  $y(0) = 0, y(0.2) = 0.02, y(0.4) = 0.0795, y(0.6) = 0.1762, y'(0) = 0, y'(0.2) = 0.1996, y'(0.4) = 0.3937, y'(0.6) = 0.5689$  (07 Marks)
- b. Prove that the shortest distance between two points in a plane is a straight line. (07 Marks)
- c. Find the extremal of the functional  $I = \int_{x_1}^{x_2} (y^2 + y'^2 + 2ye^x) dx$  (06 Marks)

\*\*\*\*\*